

Indian Journal of Mathematics

Volume 35, No. 2, 1993

CONTENTS

S. A. Settu

APPROXIMATION OF CONTINUOUS FUNCTIONS WITH GIVEN MODULUS OF
CONTINUITY BY $[F, d_n]$ MEANS 99-112

Abstract: A. A. Efimov [3] obtained uniform approximation for a class of continuous functions by Euler $(E, 1)$ means of their Fourier series. The $[F, d_n]$ methods were introduced by A. Jakimovski [5] as a generalization of both the Euler method and the Stirling-karamata-Lototsky method [6]. In this paper by considering $[F, d_n]$ methods, while we obtain new results for some known methods of $[F, d_n]$ family such as Karamata etc., we also extend the results to a larger class of continuous functions.

P. N. Natarajan

ON A GENERALIZED PRODUCT OF CERTAIN SEQUENCES OVER
NON-ARCHIMEDEAN FIELDS 113-120

Abstract: Throughout this paper sequences and infinite matrices have their entries in a complete, non-trivially valued non-archimedean field K . In this paper a generalized product of sequences is introduced and characterizations for $l_\infty(p), c_0(p)$ to be closed under this product are studied.

B. Mond and J. E. Pecaric

CONVEX INEQUALITIES FOR SEVERAL SELF-ADJOINT OPERATORS ON
A HILBERT SPACE 121-135

S. Elumalai and R. Ravi

ON REMOTAL POINTS OF PAIRS OF SETS IN NORMED LINEAR SPACES 137-146

Abstract: We give a characterization of remotal points of pairs of bounded sets in terms of extremal points. Some applications of remotal points of pairs of sets in the spaces $C(Q)$ and $C_R(Q)$ (Q -compact) are also provided.

M. M. Parmenter

ISOMORPHIC INCIDENCE ALGEBRAS OF GRAPHS 147-153

Abstract: Let $I(G, R)$ denote the incidence algebra of a locally finite directed graph G over a commutative ring R with identity. In this paper, we show that under various sets of conditions on G and R , the incidence algebra $I(G, R)$ determines the graph G .

Gaven J. Martin

THE COMMUTATOR SPECTRUM AND CONJUGACY CLASSES OF
A DISCRETE SUBGROUP OF $SL(2, C)$ 155-170

Abstract: In this paper we discuss the set of traces of commutators in an arbitrary discrete subgroup of $SL(2, C)$. That is we fix $f \in \Gamma$ and consider the set

$\{\text{tr}[f, g] : g \in \Gamma\}$. We show that generically this set is discrete in the plane by forming a series, related to the Poincaré series, which we show to converge in a half space (modulo the obvious infinite multiplicities). We then apply this result to aspects of theory of discrete groups. In particular, the finiteness of certain conjugacy classes of subgroups and geometric inequalities, such as Jørgensen's inequality.

S. S. Bhatiya and B. Ram

ON L^1 -CONVERGENCE OF CERTAIN MODIFIED TRIGONOMETRIC SUMS 171-176

Abstract: L^1 -convergence of r -th differential of trigonometric sums introduced by Ram and Kumari [3] has been studied. Results of Ram and Kumari have been obtained as corollary of our result.

R. K. Saxena and Y. Singh

INTEGRAL OPERATORS INVOLVING GENERALIZED H -FUNCTION 177-188

Abstract: The object of the present note is to introduce two new fractional integration operators associated with l -function due to Saxena [15] and to establish certain important properties which generalize the results given earlier by several authors [3, 6, 7, 9-14]. Inversion formulae and compositions (or product) of these operators are also established.
