

Indian Journal of Mathematics

(*B. N. Prasad Birth Centenary Commemoration Volume-I*)

Volume 41, No. 1, 1999

CONTENTS

- U. N. Singh**
PROFESSOR B. N. PRASAD i-xviii
- Todor Gramchev and Luigi Rodino**
GEVREY SOLVABILITY FOR SEMILINEAR PARTIAL DIFFERENTIAL EQUATIONS 1-13
Abstract: Using the fixed point theorem and the methods of the microlocal analysis, we prove a result of Gevrey solvability for a second order semilinear partial differential equation with double characteristics. As introduction, we give a quick survey on the problem of the local solvability for linear and nonlinear equations.
- Horst Herrlich and David Vaughan**
WEAK SEPARATEDNESS FOR NEARNESS SPACES 15-24
Abstract: We introduce the concept of weak separatedness for nearness spaces, a property which is strictly weaker than separatedness, but which behaves similarly. Weak separatedness is bireflective in *NEAR* and is preserved by both strict and simple completions; for weakly separated T_1 nearness spaces, the simple completion is an epireflection.
- G. Isac, V. M. Sehgal and S. P. Singh**
AN ALTERNATE VERSION OF A VARIATIONAL INEQUALITY 25-31
- Hans-Peter A. Künzi**
PERFECT PREIMAGES HAVING G_δ -DIAGONALS OF QUASI-METRIZABLE SPACES 33-37
Abstract: We show that a perfect preimage having a G_δ -diagonal of a regular quasi-metrizable space need not be quasi-metrizable.
- S. Bhargava, Chandrashekar Adiga and D. D. Somashekara**
RAMANUJAN'S ${}_1\psi_1$ SUMMATION AND AN INTERESTING GENERALIZATION OF AN IDENTITY OF JACOBI 39-42
Abstract: As a limiting case of Ramanujan's ${}_1\psi_1$ summation we obtain a generalization of Jacobi's expansion for $(q)_\infty^3$.

Valentin Gregori and Salvador Romaguera

COMMON FIXED POINT THEOREMS FOR PAIRS OF FUZZY MAPPINGS 43-54

Abstract: We prove fixed point theorems for pairs of fuzzy mappings satisfying contractive-type conditions and a rational inequality, respectively, in left \mathbf{K} -sequentially complete quasi-pseudo-metric spaces.

Sigit Nugroho and Z. Govindarajulu

TESTS FOR RANDOM EFFECTS IN TWO-WAY EXPERIMENT WITH ONE OBSERVATION PER CELL 55-66

Abstract: In this paper, we study the random effects model of the two-way experiments without interaction with one observation per cell nonparametrically. We derive the locally most powerful rank test (LMP rank test) for the random effects. We also specialize the test statistic and obtain Ψ_L when errors conform to a logistic distribution. The Ψ_L and Friedman statistic for testing no fixed treatment effects in two-way lay-out based on ranks are linearly related. The test statistic Ψ_L is discussed further by finding the asymptotic distribution of the related test under the null hypothesis and comparing with other statistics (Anderson and Fisher) for testing the hypothesis of no random treatment effects in terms of their powers when data come from normal populations.

Sehie Park

GENERALIZED BIRKHOFF-KELLOGG TYPE THEOREMS AND APPLICATIONS 67-77

Abstract: We obtain very general Birkhoff-Kellogg type theorems on eigenvectors of broad classes of compact multimaps which appear in nonlinear analysis and algebraic topology. Then are applied to fixed point and best approximation problems on such classes of multimaps defined on spheres of normed vector spaces.

J. Pečarić, G. Allasia and C. Giordano

CONVEXITY AND THE GAMMA FUNCTION 79-93

Abstract: Convexity and n -convexity properties of some functions related to the logarithm of the gamma function are proved and pertinent inequalities obtained. These results are then applied to generalize well-known inequalities.

E. Thandapani and M. Maria Susai Manuel

ON SOME CLASSES OF SOLUTIONS OF A NONLINEAR SECOND ORDER DIFFERENCE EQUATION 95-113

Abstract: The authors consider the difference equation

$$\Delta(a_n \Delta x_n) + q_n f(x_n + 1) = 0, n = 0, 1, 2, \dots \quad (*)$$

where $a_n > 0, q_n > 0$ for all $n \geq 0$ and $f : \mathbf{R} \rightarrow \mathbf{R}$ is continuous such that $uf(u) > 0$ for $u \neq 0$. Dividing the solutions of (*) into several classes, the authors obtain conditions for the existence and non-existence of solutions in these classes.

K. R. Nagarajan and T. Soundararajan

ON THE MODULE OF I-FORMS ON A DIFFERENTIABLE MANIFOLD
AND PARALLELIZABLE MANIFOLDS

115-129

Abstract: Let $A^1(M)$ be the module over $C^\infty(M)$ of all smooth 1- forms on a differentiable manifold M . Then the following conditions are equivalent: (1) $A^1(M)$ is countably generated (2) $A^1(M)$ is finitely generated (3) M is paracompact. It is also proved that M will be parallelizable if and only if $A^1(M)$ is a free module.
